



PreKUnil[®]
LNAA TABLETS[®]

The original LNAA formulation with two decades of clinical validation.

Phenylalanine free, Large Neutral Amino Acid (LNAA) tablets that allow an individual under medical supervision with proven PKU, to follow a more relaxed diet plan.

Clinically Proven

- To inhibit phenylalanine transport across blood-brain barrier¹

Indications

- For individuals (with PKU only) 8 years of age and older
- PreKUnil can also be used by untreated or late-diagnosed PKU patients

Benefits

- PreKUnil tablets relaxes the diet making it easier to manage
- Allows greater intake of natural food
- Reduces the expenses of modified low protein foods
- PreKUnil tablets are tasteless and odorless unlike the sensory characteristics of traditional formulas for PKU
- A PKU diet plan with PreKUnil allows for unlimited fruit and vegetables, more grains and cereals and meat, poultry or fish in small quantities (PreKUnil does not allow for unlimited use of high protein foods)

Warning

- Not to be used while planning a pregnancy or during a pregnancy



Products designed by healthcare professionals

solace
NUTRITION

Nutritional Information

Nutrients	Per Tablet
Calories, (kJ)	2.2, (9.11)
Protein Equivalent, g	0.4
Amino Acids, g	0.5
Carbohydrate, g	0.1
Fat, g	0.01
Trans, g	0
Vitamin B ₆ , mg	0.05
Vitamin B ₁₂ , ug	0.09

Ingredients: L-Tyrosine, dicalcium phosphate, corn starch, L- Tryptophan, L-Methionine, L-Leucine, L-Lysine, potato starch, talcum, L-Isoleucine, L-Valine, L-Threonine, L-Arginine, L-Histidine, hydroxypropylmethylcellulose, titanium dioxide, povidone, colloidal silicon dioxide, magnesium stearate, shellac, glycerol, vitamin B₆, vitamin B₁₂.

Suggested Intake: 0.5 tablet per kg of body weight per day (0.5 tablet/kg/d).

Daily intake to be determined by your healthcare professional specialized in metabolic disorders. The dosage is dependant on the age, body weight, medical condition and individual Phe tolerance. PreKUnil is not suitable as a sole source of nutrition.

Directions for Use: Use under medical supervision. Total number of tablets must be taken 3 or more times per day with meals. Always take tablets with food and at least a glass (8 fl. oz.) of water.

Example

Patient with a body weight of 60 kg

60 kg x 0.5 tablet = 30 tablets/day

Take 10 tablets 3 times a day or 7-8 tablets 4 times per day.

The tablets may be consumed during the beginning, middle or end of a meal.

PreKUnil must be supplemented with a moderately restricted PKU diet containing other sources of protein, vitamins and minerals.

An individualized diet plan must be followed at all times.

References

1) Rex A. Moats, Ph.D., Kathryn D. Moseley, M.S. R.D, Richard Koch, M.D., Marvin Nelson, M.D. Brain phenylalanine concentrations in phenylketonuria: research and treatment of adults. Pediatrics 2003 Dec;112(6 Pt 2):1575-9. (see website for full listing)



Ordering PreKUnil® LNAAs® Tablets
Call (888) 876-5223

Reimbursement code (NDC)*: 57771-0001-42

*National Drug Codes (NDC) Format Codes: Products sold by Solace Nutrition are classified as Medical Foods by the US Food and Drug Administration (FDA). NDC format codes are product codes adjusted according to standard industry practice to meet the format requirements of pharmacy and health insurance computer systems.

solace
NUTRITION

One Research Court, Suite 450
Rockville, MD 20850
Tel: 240 403-2635
Fax: 401 633-6066
Toll Free: 888 8-SOLACE
Email: info@solacenutrition.com
Visit us at www.solacenutrition.com