

Directions for Use

Daily intake to be determined by your healthcare professional specialized in mitochondrial diseases. May be consumed orally or via tube feeding.

1. Using the dosing spoon provided, measure the amount of Cyto-Q MAX[®] liquid required to meet your serving requirements.
2. Add the measured liquid to water, formula, or dosing syringe directly as a bolus. All or part of the serving may also be taken by mouth.

Warning

Not for the general population of consumers. Not for parenteral use. Not for use as a sole source of nutrition. Only for individuals with Mitochondrial Cytopathies or inborn errors of metabolism. Cyto-Q MAX[®] is oxidized when exposed to air and should be consumed directly after dispensing. **Use under medical supervision.**

Storage

All bottles (opened or unopened) should be kept in a cool, dry place (not in refrigerator). Reseal opened bottles and use within 3 months.

Cyto-Q MAX[®]

Unflavored concentrated Ubiquinol



Medical Food

A Medical Food for use in the dietary management of Inborn Errors of Metabolism such as Mitochondrial Cytopathies.

A concentrated, liposomal ubiquinol solution for enhanced absorption

Allergen tested at No Detectable Limit (NDL) for eggs, dairy, fish, crustacean, soy, corn, gluten, tree nuts, and peanut.

1 Bottle = 170 mL = 17,000 mg Ubiquinol
Net: 5.75 fl. oz (170 mL)

Nutrition Information

Nutrients	per 1 mL	
Calories	kcal	1.0
Carbohydrate	g	0.004
Protein	g	0
Fat	g	0.1
Trans	g	0
Vitamins		
Vitamin E, IU (mg α-T.E.)		1.0 (0.67)
Other		
Ubiquinol	mg	100

Ingredients

Purified Water, Coenzyme Q10 (Ubiquinol), Medium Chain Triglycerides, Sunflower Lecithin, Polysorbate 80, Ascorbyl Palmitate, Citric Acid, Potassium Sorbate and D-Alpha Tocopherol.

Distributed by:

Solace Nutrition, LLC
10 Alice Court
Pawcatuck, CT 06379
Tel: 888-8-SOLACE
Mon-Fri 9:00-4:00PM ET

solace
NUTRITION



8 57771 00702 4