

Directions of Use: GlyMAX™ is soluble in water and can be consumed either orally or via tube feeding. The daily intake of GlyMAX™ powder is to be determined by your physician or dietitian.

Oral or Tube Feeding

1. Measure (using the scoop provided or a gram scale) the recommended amount of GlyMAX™ powder.
2. For oral, consume powder by either sprinkling on food or into liquids, or taking directly by mouth followed with water.
3. For tube feeding, add powder in small increments to the formula (room temperature or cold) and stir well. GlyMAX™ powder and formula mixture should be used within 4 hours

Warning: Not for parental use. Not for use as a sole source of nutrition. Only for individuals with Multiple Acyl-CoA-Dehydrogenase Deficiency (MADD) and other Organic Acid Metabolism Disorders such as Isovaleric Acidemia. **Use under medical supervision.**

Storage: Unopened bottles should be kept in a cool, dry place. Reseal unopened bottles and use within three months.

Contact Solace Nutrition, LLC

888-8-SOLACE
Mon-Fri 9:00-4:00PM ET
www.solacenutrition.com



Medical Food

A Medical Food for the dietary management of Multiple Acyl-CoA-Dehydrogenase Deficiency (MADD) and other Organic Acid Metabolism Disorders such as Isovaleric Acidemia. Not for the general population of consumers.

Pure, soluble & slight sweet tasting Glycine powder.

Allergen-free for egg, milk, fish, crustacean, soy, gluten, corn, tree nuts, and peanut. Free of GMO, artificial flavorings and any preservatives.

Expiration/Lot #
See bottom of bottle

Net Weight: 225 g

Nutrition Information

Nutrients	per 100 g powder	
Energy from Glycine	kcal	400
Fat	g	0
Trans Fat	g	0
Carbohydrate	g	0
Protein	g	0
Glycine	g	100

Conversion: 1 level scoop = 3 g Glycine

Ingredients

Glycine

Distributed by

Solace Nutrition, LLC
Tel: 888-8-SOLAC

