

Directions of Use: CytoCarn® is soluble in water and can be consumed either orally or via tube feeding. The daily intake of CytoCarn® powder is to be determined by your physician or dietitian.

Oral or Tube Feeding

1. Measure (using the scoop provided or a gram scale) the recommended amount of CytoCarn® powder.
2. For oral, consume powder by either sprinkling on food or into liquids, or taking directly by mouth followed with water.
3. For tube feeding, add powder in 1 g (one scoop) increments to the formula (room temperature or cold) and stir well. Stir or agitate tube feeding mixture if CytoCarn® separates from formula during feeding. CytoCarn® powder and formula mixture should be used within 4 hours.

Warning: Not for parental use. Not for use as a sole source of nutrition. Only for individuals with Inborn Errors of Metabolism. **Use under medical supervision.**

Storage: Unopened bottles should be kept in a cool, dry place. Reseal unopened bottles and use within three months.

Distributed by Solace Nutrition, LLC,
10 Alice Court, Pawcatuck, CT 06379
Tel: 888-8-SOLACE Mon-Fri 9:00-4:00PM ET
www.solacenutrition.com



CytoCarn®

Medical Food

A Medical Food for Inborn Errors of Metabolism, such as Mitochondrial Cytophathies and Fatty Acid Metabolism Disorders. Not for the general population of consumers.

Pure & soluble Acetyl-L-Carnitine powder for superior absorption.

Allergen tested at No Detectable Limit (NDL) for egg, milk, fish, crustacean, soy, corn, gluten, tree nuts, and peanut. Free of GMO, animal products, artificial flavorings, and any preservatives.

Expiration/Lot #
See bottom of bottle

Net Weight: 100 g

Nutrition Information

| Nutrients | per 10 g powder | |
|--------------------|-----------------|----|
| Energy | kcal | 0 |
| Fat | g | 0 |
| Trans fat | g | 0 |
| Carbohydrate | g | 0 |
| Protein | g | 0 |
| Acetyl-L-carnitine | g | 10 |

1 level scoop = 1 g Acetyl-L-Carnitine
1 level scoop = 800 milligrams of L-Carnitine

