Directions for Use

Use under medical supervision. Daily intake to be determined by your healthcare professional specialized in mitochondrial disorders.

Each milliliter (mL) contains 100 mg of available Ubiquinol.

- Using the dosing spoon provided, measure the amount of Cyto-Q MAX liquid required to meet your serving requirements.
- Take directly by mouth or add Cyto-Q MAX liquid to any beverage, stir well and then consume.

Warning

Not for parenteral use. Not for use as a sole source of nutrition. Only for individuals with Mitochondrial Cytopathies or inborn errors of metabolism. Cyto-Q is oxidized when exposed to air and should be consumed directly after dispensing. Use under medical supervision.

Storage

All bottles (opened or unopened) should be kept in a cool, dry place (not in refrigerator). Reseal opened bottles and use within 3 months.

Distributed by:

Solace Nutrition, LLC 10 Alice Court Pawcatuck, CT 06379 Tel: (877) 872-9270

Tel: (877) 872-9270 (Mon-Fri 9:00-4:00PM ET) www.solacenutrition.com





Medical Food

A medical food for use in the dietary management of Mitochondrial Cytopathies. Not for the general population of consumers.

A concentrated, liposomal ubiquinol solution for enhanced absorption Allergen tested at No Detectable Limit (NDL) for eggs, milk, fish, crustacean, soy, corn, gluten, tree nuts, and peanut. Tested to contain non-GMO ingredients.

Net: 170 mL 1 Bottle = 170 mL = 17,000 mg Ubiquinol

Nutrition Information		
Nutrients		per 1 mL
Calories	kcal	1.0
Carbohydrate	g	0.004
Protein	g	0
Fat	g	0.1
Trans	g	0
Vitamins		
Vitamin E, IU (mg α-T.E.)		1.0 (0.67)
Other		
Ubiquinol	mg	100

Ingredients

Purified Water, Coenzyme Q10 (Ubiquinol), Medium Chain Triglycerides, Soy Lecithin, Polysorbate 80, Ascorbyl Palmitate, Citric Acid, Potassium Sorbate and D-Alpha Tocopherol.



8 || 57771 || 00702 || 1

Rev. 20190523