

**Directions for Use:** Cyto® C is soluble in water and may be consumed either orally or via tube feeding. The daily intake of Cyto® C powder is to be determined by your physician or dietitian.

#### Oral or Tube Feeding


1. Measure (using the scoop provided or a gram scale) the recommended amount of Cyto® C powder
2. For oral, consume powder by mixing into moist foods or liquids
3. For tube feeding, add powder in 1.1 g (one scoop) increments to formula (room temperature or cold) and stir well. Stir or agitate tube feeding mixture if Cyto® C separates from liquid during feeding. Cyto® C powder and liquid mixture should be used within 4 hours

**Warning:** Not for parenteral use. Not for use as the sole source of nutrition and not for the general population. Only for individuals with Mitochondrial Cytopathies. **Use under medical supervision.**

**Storage:** Unopened bottles should be kept in a cool, dry place. Once opened, reseal bottle and use within 3 months.



**Cyto**®  
*Unflavored,  
Concentrated  
Powder*



### Medical Food

A Medical Food for the dietary management of Mitochondrial Cytopathies or other Inborn Errors of Metabolism.

Suitable for a vegan diet, free of GMOs, artificial flavorings, preservatives, colors, and sweeteners.

Allergen tested at No Detectable Limit (NDL) for egg, dairy, fish, crustacean, soy, corn, gluten, tree nuts, and peanut.

Expiration/Lot #  
See bottom of container

Net Weight:  
3.53 oz (100 g)

### Nutrition Information

Nutrients	Per 1g of powder	
<b>Energy</b>	4	kcal
<b>Carbohydrate</b>	<1	g
Dietary Fiber	<1	g
<b>Vitamins</b>		
Vitamin C (L-Ascorbic Acid)	500	mg

1 level scoop = 1.1 g powder = 550 mg Vitamin C

**Ingredients:** Vitamin C (L-ascorbic acid), Phospholipids, Fatty Acids, Citrus Bioflavonoids, Maltodextrin, Citric Acid, Sodium Benzoate, Potassium Sorbate, Sucrose

#### Manufactured by:

Solace Nutrition, LLC  
10 Alice Court, Pawcatuck, CT 06379  
Tel: 888-8-SOLACE  
Mon-Fri 9:00-4:00PM ET

**solace**  
NUTRITION